




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07D 405/12, A01N 43/40 // (C07D 405/12, 319/00, 213/00) (C07D 405/12, 317/00, 213/00)	A1	(11) International Publication Number: WO 92/07848 (43) International Publication Date: 14 May 1992 (14.05.92)
(21) International Application Number: PCT/US91/07801 (22) International Filing Date: 22 October 1991 (22.10.91) (30) Priority data: 603,969 26 October 1990 (26.10.90) US (71) Applicants: UNIROYAL CHEMICAL COMPANY, INC. [US/US]; World Headquarters, Middlebury, CT 06749 (US). UNIROYAL CHEMICAL LTD./UNIROYAL CHEMICAL LTEE [CA/CA]; 25 Erb Street, Elmira, Ontario N3B 3A3 (CA). (72) Inventors: FRIEDLANDER, Barry, T. ; 245 Bristol Street, Guelph, Ontario N1H 3M3 (US). DAVIS, Robert, A. ; 1100 Notch Road, Cheshire, CT 06410 (US).		(74) Agents: THOMPSON, Raymond, D. et al.; Uniroyal Chemical Company, Inc., World Headquarters, Mid- dlebury, CT 06749 (US). (81) Designated States: AT (European patent), AU, BE (Euro- pean patent), BG, BR, CA, CH (European patent), CS, DE (European patent), DK (European patent), ES (Eu- ropean patent), FI, FR (European patent), GB (Euro- pean patent), GR (European patent), HU, IT (European patent), JP, KR, LU (European patent), NL (European patent), PL, RO, SE (European patent), SU⁺. Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>

(54) Title: 2-(3-OXYPYRIDINYL)-1,3-DIOXOLANE AND DIOXANE COMPOUNDS, COMPOSITIONS CONTAINING SAME AND PROCESS FOR CONTROLLING FUNGI USING SAME

**(57) Abstract**

A compound having structural formula (I), where R and R¹ are the same or different and are hydrogen or C₁-C₆ alkyl; R² is hydrogen, chlorine or bromine; R³ is hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, trihalomethyl, nitro, cyano, phenyl, phenyl substituted with C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, trihalomethyl, nitro or cyano, phenoxy or phenoxy substituted with C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, trihalomethyl, nitro or cyano; and n is 0 or 1 is disclosed. A composition which comprises a fungicidally effective amount of the compound and a carrier therefor is also set forth. In addition, a process for controlling phytopathogenic fungi wherein a fungicidally effective amount of the compound is applied to the locus to be protected is described.

+ DESIGNATIONS OF "SU"

Any designation of "SU" has effect in the Russian Federation. It is not yet known whether any such designation has effect in other States of the former Soviet Union.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland			SN	Senegal
CI	Côte d'Ivoire	KR	Republic of Korea	SU+	Soviet Union
CM	Cameroon	LI	Liechtenstein	TD	Chad
CS	Czechoslovakia	LK	Sri Lanka	TC	Togo
DE	Germany	LU	Luxembourg	US	United States of America
DK	Denmark	MC	Monaco		

1 2-(3-OXYPYRIDINYL)-1,3-DIOXOLANE AND DIOXANE COMPOUNDS,
 COMPOSITIONS CONTAINING SAME AND PROCESS FOR CONTROLLING
 FUNGI USING SAME

5 BACKGROUND OF THE DISCLOSURE

1. Field of the Invention

 The present invention is directed to a new class of
substituted 2-(3-oxypyridinyl)-1,3-dioxolane and -1,3-dioxane
10 compounds. More specifically, the present invention is
directed to a new class of the above noted compounds having
particular utility as fungicides.

 The control of phytopathogenic fungi is of great
economic importance not only because fungal growth on plants
15 or on parts of plants such as fruit, blossoms, foliage,
stems, tubers, roots and the like inhibit plant production,
including the plant's commercially significant portions, such
as foliage, fruit and seed, but, in addition, because fungi
reduce the overall quality of the plant crop.

20 To control or at least reduce the detrimental
effects of fungi, plants have long been treated with
fungicides. However, the enormous economic toll taken by
known fungi, as well as the continuing development of new
fungus strains resistant to known fungicides, establishes a
25 continuing need to develop new and more effective fungicides
which possess curative, preventative and systemic action to
protect all manner of plants. In addition, these newly
developed fungicides must not only provide the
above-discussed protection against the detrimental effects of
30 fungi but, negatively, must not possess properties which have
adverse effects on the plants to be protected. Furthermore,
new fungicides must not have adverse effects on the

35

SUBSTITUTE SHEET

1 commercial exploitation of the crop of the plant protect d.
Finally, the fungicide must be free of characteristics which
adversely affect the surrounding environment including the
soil and the surroundings into which the fungicide is
5 introduced.

The above remarks establish the need in the art for
new compounds, distinguished from the compounds utilized in
the prior art, that provide more effective fungicidal
activity, without adverse effects on the plants to be
10 protected or the environment in which the plants are grown,
against the scourge of phytopathogenic fungi.

2. Background of the Prior Art

The prior art includes teachings directed to the
15 use of oxypyridinium compounds that are useful as fungicides.
One such reference is U.S. Patent 4,143,144, issued to Tobol
et al., which discloses certain 2-alkoxy- and 2,6-dialkoxy-4-
halomethylpyridine compounds which find utility as
fungicides. It is emphasized, however, that although the
20 compounds of the '144 patent are oxypyridinium compounds,
they have no relation to any class of substituted 1,3-
dioxolanes or substituted 1,3-dioxanes.


Similarly, U.S. Patent 4,678,504 to Schulz et al.
sets forth another class of oxypyridinium compounds useful as
25 fungicides. The compounds of Schultz et al. are salts of
O-substituted 3-oxypyridinium compounds. More specifically,
Schulz et al. describes benzyl bromide salts of
3-tetradecyloxy pyridine, 2,4'-difluorobenzylhydroxypyridine
and the like. This class of salts, like the compounds of the
30 Tobol et al. patent, teach the use of oxypyridinium compounds
as fungicidal agents. However, there is no disclosure,
suggestion or even hint of utilizing 1,3-dioxolane and
1,3-dioxane compounds which include substituted and
unsubstituted 2-(3-oxypyridinyl) substituents.

35

1 BRIEF SUMMARY OF THE INVENTION

A new class of compounds have now been developed which not only provides excellent protection against the ravages of phytopathogenic fungi but does so without
 5 adversely affecting the plants to be protected or significantly altering the environmental conditions of the soil and the surroundings into which these fungicides are introduced.

In accordance with the present invention a new
 10 class of compounds having the structural formula



where R and R¹ are the same or different and are hydrogen or C₁-C₆ alkyl; R² is hydrogen, chlorine or bromine; R³ is
 20 hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, trihalomethyl, nitro, cyano, phenyl, phenyl substituted with C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, trihalomethyl, nitro, cyano, phenoxy or phenoxy substituted with C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, trihalomethyl, nitro or cyano; and n is 0 or 1 is
 25 disclosed.


In addition, a composition comprising a fungicidally effective amount of the compound of this invention and a carrier therefor is set forth.

Finally, the present invention is also directed to
 30 a process for controlling phytopathogenic fungi which comprises applying a fungicidally effective amount of the compound of this invention to the locus to be protected.

1 DETAILED DESCRIPTION

The compounds of the present invention have the structural formula


5



- 10 where R and R¹ are the same or different and are hydrogen or C₁-C₆ alkyl; R² is hydrogen, chlorine or bromine; R³ is hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, trihalomethyl, nitro, cyano, phenyl, phenyl substituted with C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, trihalomethyl, nitro or cyano, phenoxy
 15 or phenoxy substituted with C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, trihalomethyl, nitro or cyano; and n is 0 or 1.

- It is emphasized that the compound having the structural formula I encompasses dioxolanes, which correspond to the compound having the structural formula I where n is 0.
 20 The dioxolane compounds within the contemplation of the present invention have the structural formula


25



- 30 where R, R¹, R² and R³ have the meanings given for the compound having the structural formula I.

- The equally preferred dioxanes, within the contemplation of the present invention, are defined by the compound having the structural formula I where n is 1. The
 35 dioxanes of the present application have the structural formula

SUBSTITUTE SHEET



where R, R¹, R² and R³ have the meanings given for the compound having the structural formula I.

The present invention also encompasses salts of the compound having the structural formula I. Salts within the contemplation of the subject invention are preferably addition products of the compound having the structural formula I, as well as compounds of structural formulae II and III within the genus of the compound having the structural formula I, and various acids. Among the preferred acids, which may be employed to produce salts of the present invention, are such inorganic acids as hydrochloric, hydrobromic, nitric and sulfuric acids. Organic acids, which also may be employed in the formation of salts within the contemplation of the present invention, include the sulfonic acids, methanesulfonic acid and p-toluenesulfonic acid. Of these acids, hydrochloric acid and methanesulfonic acid are particularly preferred in producing the salts of the subject invention.


A careful review of the structural formula I makes apparent to those skilled in the art that when R and R¹ are different, the carbon atom at the 2-position is asymmetrical. As such, this gives rise to optical isomers. Furthermore, in the case where either R or R¹ are not hydrogen, compounds having the structural formula I, or the subgeneric structural formulae II or III, may exist as geometric isomers in that R or R¹ are cis or trans relative to the substituents at the 2-position of the dioxolane or dioxane ring. It should therefore be appreciated that isomers, as well as mixtures of isomers, of the sort discussed above, are within the contemplation of the present invention.

SUBSTITUTE SHEET


1 Preferably, compounds having the structural formula
 I are characterized by R and R¹ being the same or different
 and being hydrogen or C₁-C₄ alkyl; R² being hydrogen or
 chlorine; R³ being hydrogen, halogen, phenyl or phenoxy
 5 substituted with methyl, ethoxy, halogen or trifluoromethyl;
 and n is 0 or 1.

More preferably, the compound having the structural
 formula I is defined by R and R¹ being hydrogen or C₁-C₃
 alkyl; R² being hydrogen, R³ being chlorine, phenyl, or
 10 phenoxy substituted with halogen or trifluoromethyl; and n is
 0 or 1. The latter definition, in the more preferred
 definition of the compound of the present invention, the
 definition of n being 0 or 1, emphasizes that no preference
 should be given over each other to the compounds having the
 15 structural formulae II and III.

The compound having the structural formula I is
 prepared in a two-step process wherein an alpha-haloketone
 having the structural formula



where X is halogen, preferably, bromine; and R³ has the
 25 meanings given in the definition of structural formula I, is
 subjected to cyclization by reaction with a diol having the
 structural formula




where R, R¹ and n have the meanings given in defining
 structural formula I.

35


SUBSTITUTE SHEET

The product of this cyclization reaction is a 2-halomethyl ketal having the structural formula



where R, R¹, R³, X and n have the meanings given above. It is emphasized that details of this type of reaction are known in the art. For example, Synthesis, 1, 23, (1974) provides specifics of this reaction. The above referenced article is incorporated herein by reference.

The 2-halomethyl ketal compound having the structural formula VI is reacted with a 3-hydroxypyridine compound having the structural formula



where R² has the same meanings as given for that substituent in structural formula I. This reaction occurs in the presence of an acid acceptor, preferably potassium carbonate. Moreover, the 3-hydroxypyridine compound having the structural formula VII is introduced into the reaction as a solution in which the solvent is preferably N,N-dimethylacetamide. The reaction usually occurs by refluxing the compounds having the structural formulae VI and VII at the boiling temperature of the solvent, usually N,N-dimethylacetamide, for a period of between about 6 hours and about 72 hours, preferably about 18 hours. The product of this reaction is the compound having the structural formula I.

SUBSTITUTE SHEET

1 Another aspect of the present invention is a
composition which comprises a fungicidally effective amount
of the compound having the structural formula I and a carrier
therefor. Carriers within the contemplation of the
5 composition of this invention may be liquids, solids or
mixtures thereof.

Turning first to liquid carriers useful in
producing the composition of the present invention, the
liquid carrier may be a solvent or a dispersant. In
10 addition, two liquid carriers can be utilized, one
serving as a solvent and the other as a dispersant.

In the preferred embodiment wherein the composition
is a solution, the solvent carrier is usually an organic
compound which may be polar or non-polar. Solvents within
15 the contemplation of the composition of this invention
include acetone, methanol, isopropanol, t-butyl alcohol,
cyclohexanone, n-butyl alcohol, toluene, xylene, dioxane,
dimethylformamide, dimethylsulfoxide, ethylene dichloride and
N-methylpyrrolidone.

20 Another class of liquid compositions within the
contemplation of this invention is emulsions. An emulsion is
formed when the compound having the structural formula I is
dispersed in water in the presence of a surface active agent.
An emulsion is preferably formed by first preparing a
25 solution of the type discussed in the above paragraph. The
solution is then dispersed in water and a surface active
agent added thereto to form the emulsion. Surface active
agents suitable for use in forming an emulsion within the
contemplation of the composition of this invention are known
30 to those skilled in the art. McCutcheon's Detergents and
Emulsifiers, Allured Publishing Corp., Ridgewood, NJ (1970);
U.S. Patent 2,514,916 and U.S. Patent 2,547,734 provide
examples of surface active agents useful in forming emulsions

35

SUBSTITUTE SHEET

1 within the contemplation of the composition of this
invention. The above recited references are incorporated
herein by reference. As set forth in these references, the
surface active agents may be anionic, cationic or non-ionic.

5 A third class of liquid compositions within the
scope of this invention utilize a liquid dispersant as the
carrier. In this embodiment, the compound having the
structural formula I is dispersed in water in the absence of
a surface active agent. Alternatively, the liquid
10 composition involves a solution of the compound having the
structural formula I, which, in turn, is dispersed in water,
again in the absence of a surface action agent.

Yet another class of liquid compositions within the
scope of the present invention utilizes an aerosol. An
15 aerosol is liquid under pressure but is gaseous at
atmospheric pressure and ambient temperature. In most
instances an aerosol composition is prepared by first forming
a solution of the compound having the structural formula I in
a conventional solvent of the type discussed above. This
20 solution is admixed with a volatile liquid aerosol under
pressure in which condition the composition is applied.

A second major class of composition within the
meaning of the present invention are solid compositions which
employ a solid carrier. Solid carriers useful in the
25 formation of the composition of this invention include dusts,
granules, wettable powders, pastes and water soluble solids.
For example, compositions within the contemplation of this
invention may be applied as a dust when adsorbed or absorbed
onto or mixed with a powdered, solid carrier. A solid
30 carrier such as a mineral silicate, e.g., mica, talc,
pyrophyllite and clays, may be utilized for this purpose.

35

SUBSTITUTE SHEET

1 Additional solid compositions can be prepared from
granular formulations of the compound having the structural
formula I and a granular or pelletized form of carrier such
as granular clay, vermiculite, charcoal, corncobs or the
5 like. The use of granular formulations is particularly
suitable for application by broadcasting, side-dressing, soil
incorporation or seed treatment.

A mixture of a solid and liquid composition, that
employs both a liquid and a solid carrier, may also be used.
10 Such a composition, for example, is prepared by dispersing a
solid, on which the compound having the structural formula I
is absorbed or adsorbed, in a liquid dispersant. Such a
composition preferably includes a surface active agent to
maintain the solid particles dispersed in the liquid
15 dispersant.

It should be emphasized that the composition of the
present invention may utilize a carrier which is itself
active. That is, the carrier may be a plant growth regulant,
an insecticide, an acaricide, a fungicide, a bactericide or
20 the like.

The concentration of the compound having the
structural formula I in the composition of this invention is
a fungicidally effective amount. The exact concentration of
a fungicidally effective amount depends upon such factors as
25 the specific plant, or plants, which is to be protected, the
fungus, or the fungi, which is to be controlled, soil
conditions and chemistry and the climatic conditions under
which the plant is grown. Generally, the concentration of
Compound I, which is representative of a fungicidally
30 effective amount, in the composition of the present invention
may range from about 0.1% to about 95% by weight. However,
when the compound having the structural formula I is applied

35

SUBSTITUTE SHEET

1 as a spray, the dilution may be very high, resulting in a
concentration as low as a few parts per million parts of
composition. On the other hand, when ultra-low volume
applications are employed, full strength concentrates may be
5 utilized.

The present invention is also concerned with a
process for controlling phytopathogenic fungi. In this
process the compound having the structural formula I is
applied to the locus under attack by said fungi in a
10 fungicidally effective concentration.

In one preferred embodiment of the process of this
invention the compound having the structural formula I is
applied to the foliage of the plant or plants to be
protected. This so-called "foliar treatment" is effectuated
15 by applying the compound having the structural formula I to
foliage at a concentration of between about 10 milligrams and
about 500 milligrams of the compound per liter of inert
liquid carrier.

In another preferred embodiment of the process of
20 the present application, the process of controlling
phytopathogenic fungi, a fungicidally effective amount of the
compound having the structural formula I is applied to the
soil in which the plant, or plants, to be protected from
fungi is grown. In this method, the so-called "systemic
25 treatment," the compound having the structural formula I is
applied to the soil in a concentration of between about 0.125
and about 10 kilograms of compound per hectare of soil in
which the plant, or plants, to be protected is grown. More
preferably, systemic control involves application of between
30 about 0.125 kg/ha and about 5 kg/ha of the compound having
the structural formula I to the soil in which the plant, or
plants, to be protected is grown.

35

SUBSTITUTE SHEET.

1 Independent of which preferred process of
controlling fungi is utilized, the application may be applied
prior to or after infection by fungi. Furthermore, it should
be appreciated that the exact dosage, applied systemically or
5 directly to foliage, is dictated by the fungus to be
controlled and the particular plant to be protected.

In another embodiment of the process of controlling
phytopathogenic fungi utilizing the compound of the present
invention, the compound having the structural formula I, the
10 compound is applied as a coating to seeds of the plant to be
protected. This embodiment of the process of the present
invention provides the benefits of the two preferred
embodiments discussed above, foliage treatment and systemic
treatment. That is, the fungicidal coating, the coating of
15 the compound having the structural formula I, protects the
soil from infection by the fungi but is also taken up by the
plants systemically to protect the plant from fungal attack.
In this so-called "seed coating method," a concentration of
the compound having the structural formula I in the range of
20 between about 5 and about 75 grams per 100 kg. of seed is
usually utilized.

The following examples are given to illustrate the
scope of the present invention. Because these examples are
given for illustrative purposes only, the scope of the
25 instant invention should not be limited thereto.

30

35

SUBSTITUTE SHEET

EXAMPLE 1

1

Preparation of 3-[[2-([1,1'-Biphenyl]-4-yl)-1,3-dioxolan-2-yl]methoxy]pyridine (Compound No. 2)

5

A mixture of 2-([1,1'-biphenyl]-4-yl)-2-(bromo-methyl)-1,3-dioxolane (16 g.), 3-hydroxypyridine (5.7 g.) and potassium carbonate (9 g.) were refluxed with stirring in N,N-dimethylacetamide (75 ml.) for 18 hours. After refluxing, the mixture was allowed to cool and was thereafter poured into water (150 ml.). The resultant aqueous solution was extracted twice with ether. The combined ether fractions were washed once with 5% aqueous sodium hydroxide and 4 times with water. The washed ether fractions were then dried, filtered and the solvent evaporated to yield 3-[[2-([1,1'-biphenyl]-4-yl)-1,3-dioxolan-2-yl]methoxy]pyridine as an oil.

20

25

30

35

SUBSTITUTE SHEET

1

EXAMPLE 2

Preparation of 3-[[2-([1,1'-Biphenyl]-4-yl)-1,3-dioxolan-2-yl]-methoxy]pyridine hydrochloride (Compound No. 1)

5

The compound of Example 1, 3-[[2-([1,1'-biphenyl]-4-yl)-1,3-dioxolan-2-yl]methoxy]pyridine (3 g.) was formed into a solution by being dissolved in dry ether (250 ml.). The solution was cooled in an ice bath and dry hydrogen chloride gas was bubbled into the solution. This resulted in precipitation. When precipitation was ended, the solid was isolated by filtration and crushed to give a fine powder. The powder was identified as the titled compound having a melting point of 191°C to 193°C.

15

20

25

30

35

SUBSTITUTE SHEET

EXAMPLE 3Preparation of 3-[[2-(4-Bromophenyl)-1,3-dioxolan-2-yl]-methoxy]pyridine methanesulfonate (Compound No. 7)

5 A mixture of 2-(bromomethyl)-2-(4-bromophenyl)-1,3-dioxolane (25.5 g.), 3-hydroxypyridine (9.0 g.) and potassium carbonate (14.2 g.) in N,N-dimethylacetamide (125 ml.) was refluxed for 18 hours with stirring. Upon completion of this
10 refluxing the mixture was allowed to cool. To this cooled reaction mixture was added water (250 ml.) and extracted with two portions of ether. The combined ether fractions were washed once with 5% aqueous sodium hydroxide and four times with water. The resulting ether layer was dried, filtered
15 and evaporated resulting in the formation of an oil (18.5 g.).

A solution of this oil (3 g.) in dry ether was prepared and cooled in an ice bath. To this cooled solution was added methanesulfonic acid (0.9 g.) by dropwise addition accompanied by stirring. The resulting precipitate was
20 obtained by filtration. After drying, the product, having a melting point of 133°C to 134°C, was identified as 3-[[2-(4-bromophenyl)-1,3-dioxolan-2-yl]methoxy]pyridine methanesulfonate.

25

30

35

SUBSTITUTE SHEET

EXAMPLE 4

Preparation of 3-[[2-([1,1'-biphenyl]-4-yl)-4,5-dimethyl-1,3-dioxolan-2-yl]methoxy]pyridine (Compound No. 15)

A mixture of 2-([1,1'-biphenyl]-4-yl)-2-(bromo-methyl)-1,3-dioxolane (10 g.), 3-hydroxypyridine (4.7 g.) and potassium carbonate (5.2 g.) was refluxed with stirring for 18 hours. At the completion of this period, the mixture was cooled to ambient temperature. Water (250 ml.) was added to the cooled mixture and extracted with toluene. The resultant toluene solution was washed with water, then dried, filtered and evaporated to yield 3-[[2-([1,1'-biphenyl]-4-yl)-4,5-dimethyl-1,3-dioxolan-2-yl]methoxy]pyridine as an oil.

SUBSTITUTE SHEET

1

EXAMPLE 5Preparation of 3-[[2-(2,4-Dichlorophenyl)-4-methyl-1,3-dioxolan-2-yl]methoxy]pyridine (Compound No. 18)

5

A mixture of 2-(bromophenyl)-2-(2,4-dichlorophenyl)-4-methyl-1,3-dioxolane (10 g.), 3-hydroxypyridine (3.8 g.) and potassium carbonate (5.5 g.) in N,N-dimethylacetamide (90 ml.) was refluxed with stirring for 18 hours. After the mixture was allowed to cool, it was poured into water and extracted with toluene. The toluene fraction, thereby obtained, was washed with 5% aqueous sodium hydroxide and evaporated to yield 3-[[2-(2,4-dichlorophenyl)-4-methyl-1,3-dioxolan-2-yl]methoxy]pyridine as an oil.

15

20

25

30

35

SUBSTITUTE SHEET

Preparation of 3-[[2-[4-[3-(Trifluoromethyl)phenoxy]phenyl]-1,3-dioxolan-2-yl]methoxy]pyridine hydrochloride
(Compound No. 22)

A mixture of 2-(bromomethyl)-2-[4-[3-(trifluoromethyl)phenoxy]phenyl]-1,3-dioxolane (8 g.), 3-hydroxypyridine (2.4 g.) and potassium carbonate (3.5 g.) in N,N-dimethylacetamide (40 ml.) was refluxed for 18 hours with stirring. The reaction mixture was allowed to cool, poured into water and extracted twice with toluene. The combined toluene fractions were washed once with 5% aqueous sodium hydroxide and three times with water. The toluene fraction was then dried, filtered and evaporated to yield an oil (5.3 g.).

A portion of the thus obtained oil (2 g.) was dissolved in ether and dry hydrogen chloride gas was bubbled into the solution. This resulted in material oiling out of the solution. The material oiled out of the solution was cooled resulting in the solidification of the oil. This solid was isolated by filtration and identified as 3-[[2-[4-[3-(trifluoromethyl)phenoxy]phenyl]-1,3-dioxolan-2-yl]methoxy]pyridine hydrochloride. This product was obtained as a gray powder and was identified by its melting point of 138°C to 140°C.

30

35

SUBSTITUTE SHEET

EXAMPLE 7

1

Preparation of Compounds Nos. 3-6, 8-14, 16, 17, 19-21
and 23-29

5

Compound Nos. 3-6, 8-14, 16, 17, 19-21 and 23-29 were prepared in accordance with the procedure used in Examples 1 to 6. They resulted in the formation of compounds whose identity is summarized in Table I. Table I identifies not only the compounds of this example but also the compounds synthesized in Examples 1 to 6. These compounds, Compound
10 Nos. 1 to 29, are identified by their structural formula wherein the substituents are defined in Table I. Those compounds which are salts are further defined by the identity of the acid which is employed to form the salt. In addition,
15 those compounds which are solid at ambient conditions are defined by their melting point. Table II identifies those compounds listed in Table I as being an oil at ambient conditions by their nuclear magnetic resonance spectroscopy
20 data.

25


30

35

SUBSTITUTE SHEET

-20-

TABLE I



Cpd. No.	n	R	R ¹	R ²	R ³	Salt	MP(°C)
1	0	H	H	H	4-Ph	HCl	191-193
2	0	H	H	H	4-Ph		Oil
3	0	H	H	H	2,4-Cl ₂		Oil
4	0	H	H	H	4-Br		Oil
5	0	H	H	H	2-Cl		Oil
6	0	H	H	H	4-CH ₃	CH ₃ SO ₃ H	104-106
7	0	H	H	H	4-Br	CH ₃ SO ₃ H	133-134
8	0	H	H	H	4-C(CH ₃) ₃	HCl	134-137
9	0	CH ₃ CH ₂ CH ₂	H	H	H		Oil
10	0	CH ₃ CH ₂	H	H	H		Oil
11	0	CH ₃	CH ₃	H	H		Oil
12	0	CH ₃	H	H	4-Ph		Oil
13	0	CH ₃	H	H	4-Cl		Oil
14	0	CH ₃	H	H	2-Cl		Oil
15	0	CH ₃	CH ₃	H	4-Ph		Oil
16	0	H	H	H	4-F	HCl	168-172
17	0	CH ₃ CH ₂	H	H	4-Ph		Oil
18	0	CH ₃	H	H	2,4-Cl ₂		Oil
19	0	CH ₃ CH ₂	H	H	2,4-Cl ₂		Oil
20	0	H	H	H	4-(4-CH ₃ OPh)		Oil
21	0	H	H	H	4-(3-CF ₃ OPh)		Oil
22	0	H	H	H	4-(3-CF ₃ OPh)	HCl	138-140
23	0	CH ₃	H	H	4-(3-CF ₃ OPh)		Oil

SUBSTITUTE SHEET

-21-

TABLE I (Cont'd.)

<u>Cpd.</u> <u>No.</u>	<u>n</u>	<u>R</u>	<u>R¹</u>	<u>R²</u>	<u>R³</u>	<u>Salt</u>	<u>MP(°C)</u>
24	0	CH ₃	H	H	4-(2,4-Cl ₂ OPh)		Oil
25	0	H	H	H	4-(2-CH ₃ CH ₂ OOPh)		Oil
26	0	H	H	2-Br	4-(4-FOPh)	HCl	73-78
27	0	H	H	2-Cl	4-(4-BrOPh)		Oil
28	0	H	H	2-Cl	4-(4-ClOPh)		Oil
29	1	H	H	5-Cl	2,4-Cl ₂	HCl	128-129

Remarks:

Ph is phenyl

OPh is phenoxy

SUBSTITUTE SHEET

TABLE II

<u>Cpd No.</u>	<u>NMR data (δ) for CDCl₃</u>
2	8.1-8.4(m,2H), 7.0-7.7(m,11H), 4.3(s,2H), 3.7-4.3(m,4H)
3	8.1-8.4(m,2H), 7.7(d,1H), 7.0-7.5(m,4H), 4.4(s,2H), 3.7-4.3(m,4H)
4	8.1-8.3(m,2H), 7.4(s,4H), 7.1-7.3(m,2H), 4.2(s,2H), 3.8-4.2(m,4H)
5	8.1-8.4(m,2H), 7.7-7.9(m,1H), 7.1-7.4(m,5H), 4.5(s,2H) 3.8-4.3(m,4H)
9	8.1-8.4(m,2H), 7.1-7.7 (m,7H), 4.2(s+s,2H), 3.4-4.3(m,3H), 0.8-1.8(m,7H)
10	8.1-8.4(m,2H), 7.1-7.7(m,7H), 4.2(s+s,2H), 3.4-4.4 (m,3H), 1.3-1.8(m,2H), 0.8-1.2(m,3H)
11	8.1-8.4(m,2H), 7.1-7.7(m,7H), 4.2(s,2H), 3.4-4.1(m,2H), 2.3(d,3H), 1.2(d,3H)
12	8.1-8.5(m,2H), 7.1-7.7(m,11H), 4.2(s+s,2H), 3.5-4.2(m,3H), 1.3(d+d,3H)
13	8.1-8.4(m,2H), 7.1-7.7(m,6H), 4.2(s+s,2H), 3.5-4.6 (m,3H), 1.3(d+d,3H)
14	8.1-8.4(m,2H), 7.1-7.9(m,6H), 4.5(s+s,2H), 3.4-4.5(m,3H), 1.3(d+d,3H)

SUBSTITUTE SHEET

TABLE II (Cont'd.)

<u>Cpd. No.</u>	<u>NMR data (δ) for CDCl₃</u>
15	8.1-8.4(m,2H), 7.2-7.7(m,11H), 4.2(s,2H), 3.5-4.1(m,2H), 1.4(d,3H), 1.2(d,3H)
17	8.0-8.6(m,2H), 7.1-7.7(m,11H), 4.2(s+s,2H), 3.5-4.3(m,3H), 1.3-2.0(m,2H), 0.8-1.2(m,3H)
18	8.1-8.3(m,2H), 7.5-7.8(m,1H), 7.1-7.4(m,4H), 4.4(s+s,2H), 3.3-4.4(m,3H), 1.3(d+d, 3H)
19	8.1-8.3(m,2H), 7.5-7.8(m,1H), 7.1-7.5(m,4H), 4.4(s+s,2H), 3.4-4.3(m,3H), 1.3-1.9(m,2H), 0.8-1.2(m,3H)
20	6.8-7.7(m,12H), 4.3(s,2H), 3.7-4.3(m,4H), 2.3(s,3H)
21	8.0-8.7(m,2H), 6.9-7.8(m,10H), 4.3(s,2H), 3.8-4.4(m,4H)
23	8.1-8.4(m,2H), 6.9-7.7(m,10H), 4.2(s+s,2H), 3.5-4.6(m,3H), 1.3(d+d,3H)
24	8.0-8.6(m,2H), 6.8-7.7(m,9H), 3.5-4.5(m,6H), 1.3(d+d,3H)
25	8.1-8.4(m,2H), 6.8-7.6(m,10H), 4.2(s,2H), 3.9(q,2H), 3.8-4.2(m,4H), 1.2(t,3H)
27	7.9-8.1(m,1H), 6.8-7.7(m,9H), 4.2(s,2H), 3.9-4.3(m,4H)
28	7.8-8.0(m,1H), 6.8-7.7(m,9H), 4.2(s,2H), 3.8-4.4(m,4H)

SUBSTITUTE SHEET

EXAMPLE 8Preparation of Fungicidal Compositions

Each of Compound Nos. 1 to 24 (0.3 g.), summarized in Table I, were dissolved in acetone (10 ml.). Water and one or two drops of the emulsifying agent, Triton [trademark] X-100 were added to the thus formed solution to form an emulsion. The amount of water added was a function of the desired concentration of the emulsion composition which are hereinafter reported as milligrams per liter (mg/l).

SUBSTITUTE SHEET

EXAMPLE 9

1

Control of Powdery Mildew Fungus by Systemic Root Uptake

5

Compositions of Compound Nos. 1 to 29, formed in accordance with the procedure of Example 8, were tested to evaluate their effectiveness in preventing or controlling powdery mildew disease of barley caused by the fungus, Erysiphe graminis and powdery mildew disease of cucumber caused by the fungus, Erysiphe cichoracearum. This prevention or control capability, utilizing the compounds of the present invention, was tested using the systemic root uptake method of applying the fungicidal compositions to the plants to be protected.

10

In accordance with the aim of this test, pots (4 x 4 x 3.5 inches) containing ten plants of barley (Variety "Herta") or ten plants of cucumber (Variety "Marketmore 70") were grown to an age of six days and ten days, respectively. Upon reaching these ages, emulsion compositions (45 ml.) of Compound Nos. 1 to 29, formed in accordance with the procedure of Example 8, were added to each pot. That is, emulsion compositions (45 ml.) of each of the compounds of Table I was separately added to pots containing ten barley or ten cucumber plants of the type enumerated above. The emulsion compositions saturated the soil in each pot without significant loss through drainage into saucers below the pots. Each of the compositions contained one of Compound Nos. 1 to 29 in a concentration of 250 mg/l. A number of pots containing the barley and cucumber plants which were treated with the compounds of this invention were left untreated as controls.

25

30

35

The barley and cucumber plants in all the pots, including those treated and those untreated, were inoculated with powdery mildew fungus 24 hours after treatment with the emulsion compositions containing Compound Nos. 1 to 29.

SUBSTITUTE SHEET

1 Fungus inoculation was accomplished by tapping leaves of
previously infected barley and cucumber plants over the
treated and untreated pots containing the barley and cucumber
plants, respectively, to distribute the spores of the fungus
over the plants growing in the pots.

5 Six days after inoculation, disease control was
evaluated on a 0 to 6 rating scale. A 0 rating was assigned
when no disease was evidenced on the plant. A 6 rating was
given for severe disease. Intermediate ratings were assigned
depending on the degree of disease. Percent control was
10 computed by comparing the ratings of the treated and
untreated plants.

The results of this test are reported in Table III,
wherein systemic control of powdery, mildew disease in barley
is reported under the title "BMS 250." Control of powdery
15 mildew disease in cucumber is reported under the title "CMS
250."

20

25

30

35

SUBSTITUTE SHEET

EXAMPLE 10

1

Control of Powdery Mildew Fungus by Foliar Application

5 A large number of planting pots, identical with those utilized in Example 9, sufficient to accommodate testing in duplicate or triplicate for each of the 29 compounds tabulated in Table I, were each planted with eight barley plants (Variety "Larker"). There were enough planted pots so that for each treated pot there was at least one pot
10 untreated which acted as a control.

Each of the treated pots were sprayed with emulsions of each of Compound Nos. 1 to 29 in a concentration of 1000 milligrams of the compound per liter of water (1000
15 mg/l). After the foliage of the sprayed barley plants were dried, all the plants, including the unsprayed control plants, were placed in a greenhouse maintained at 21°C. All the plants, controls as well as sprayed plants, were thereupon inoculated with powdery mildew fungus, Erysiphe
20 graminis. Inoculation of the fungus was accomplished in accordance with the method utilized in Example 9.

Five days after inoculation, the plants were evaluated and assigned a disease rating of 0 to 6 in accordance with the criteria explained in Example 9. Percentage control was computed in accordance with the method
25 utilized in Example 9. The results of this test are summarized in Table III under the title "BMP 1000."

30

35

SUBSTITUTE SHEET

5

-28-

EXAMPLE 11

10

Bean Rust Fungus Eradicant and Bean Powdery
Mildew Protectant Test

Two pinto bean plants, *P. vulgaris*, were planted in a plurality of pots. When the plants were seven days old, at the primary leaf stage of growth, they were all sprayed with a suspension containing 20,000 spores of the bean rust fungus, *Uromyces phaseoli*, per milliliter of suspending water. All the pots containing the inoculated plants were than incubated in a controlled environmental chamber, maintained at 99% humidity and 21°C, for 24 hours to allow infection to develop. The plants were then moved from the incubator and allowed to dry.

Two days after inoculation the infected plants were separately sprayed with each of the emulsion compositions containing Compound Nos. 1 to 29, tabulated in Table I, prepared in accordance with the procedure of Example 8. Each of the emulsion compositions provided a dosage of 1,000 mg/l of the active compound. An equal number of infected plants, not sprayed, acted as controls. All the sprayed and unsprayed plants were placed in a greenhouse, maintained at a temperature of 21°C for five days to allow any disease to be expressed. The treated plants were additionally tested for bean powdery mildew protectant use by inoculating all plants by tapping spores of *Erysiphe polygoni* mildew over the leaves.

The sprayed and unsprayed control plants were examined for disease which was assessed using the 0 to 6 rating system described in Example 9. Disease control, as explained in Example 9, was then determined. The control of disease, expressed as percent reduction of disease, is included in Table III under the titles "BRE 1000 and PMP 1000."

SUBSTITUTE SHEET

EXAMPLE 12Control of Barley Blast by Foliar Treatment

A plurality of planting pots which included ten plants of six day old barley (Variety "Herta") were prepared. These pots were sprayed with the emulsion compositions, prepared in accordance with the procedure of Example 8, of each of Compound Nos. 1 to 29, set forth in Table I.

The plants in these pots, as well as an equal number of six day old Variety "Herta" barley plants in control pots, were inoculated with spores of the blast fungus, Pyricularia oryzae. The method of inoculation was identical with that utilized in Example 9.

All the fungus inoculated plants were placed in a greenhouse, maintained at a temperature of 21°C and a humidity of 99%, for five days. At that time, the plants were evaluated on the 0 to 6 disease rating system, explained in Example 9. Percent control was computed by comparing the ratings of the sprayed and unsprayed plants. The results of this test are included in Table III under the title "BBL 1000."

SUBSTITUTE SHEET

1

EXAMPLE 13Control of Nine Fungus Species

5 Each of the compounds included in Table I, Compound Nos. 1 to 29, were solubilized in acetone at a concentration of 500 mg/l. Filter paper discs, each 11 mm. in diameter, were immersed in each of the thus formed solutions. The discs were allowed to dry in air, thus driving off the
10 acetone solvent. An equal number of discs were not immersed in these solutions and acted as controls.

 Each of the treated and untreated disks were then placed in agar plates and the fungus species, Alternaria solani, was added to the center of each paper disc in the
15 form of a culture plug with the fungus mat in contact with the treated paper of the test disc or, in the case of the controls, in contact with the untreated filter test disc. The agar plates were incubated at 29°C in an oven.

 Upon removal from the oven, percent growth
20 inhibition by the compounds of the fungus species was evaluated by measuring the radius from the center of the fungus colony of the treated discs compared to the radius from the center of the fungus colony of the untreated disks. That is, inhibition effectuated by each of the compounds was
25 determined as a function of the percent difference between the radii of the treated and untreated discs. The results of these tests appear in Table III under the title "ALT 500."

 In a similar fashion, the fungus species: Botrytis cinerea, Fusarium oxysporum, Helminthosporium maydis,
30 Phytophthora infestans, Sclerotinia sclerotiorum and Sclerotium rolfsii were treated in exact accordance with the procedure utilized to test the fungus species, Alternaria solani. The effectiveness of Compound Nos. 1 to 29, in

35

SUBSTITUTE SHEET

-31-

5

controlling these six fungus species are summarized in Table III under the titles "BOT 400," "FUS 500," "HMA 500," "PHY 500," "SCM 500" and "SCO 500." These designations refer to the six fungus species in the order of their earlier recitation in the discussion of this example.

A separate test was utilized to determine the control of a ninth fungus species, Cercospora arachidicola. This species was tested by depositing two drops of a spore suspension of the fungus (20,000 spores per millimeter) to the above-discussed chemically treated discs rather than as a mycelial cell culture plug. Scoring of the effectiveness of the compounds in controlling the Cercospora arachidicola fungus was accomplished without the use of a control. A rating of 100 represented complete inhibition of germination and growth of the fungus; a rating of 80 represented nearly complete inhibition but some growth of the fungus; a rating of 50 represented partial inhibition but some growth or early complete inhibition with later growth; a rating 20 indicated some, but not significant, inhibition of growth; and a 0 rating indicated complete growth of the fungus without any inhibition. The rating of this species is included in Table III under the title "CER 500."

SUBSTITUTE SHEET

-32-

TABLE IIIPercent Fungicidal Control

Cpd.	ALT	BBL	BMP	BMS	BOT	BRE	CER	CMS	FUS	HMAY	PHY	PMP	SCH	SCO
No.	500	1000	1000	250	500	1000	500	250	500	500	500	1000	500	500
1	25	90	90	0	0	0	0	0	0	0	45	0	0	10
2	0	90	90	0	0	0	0	15	15	40	76	0	0	40
3	0	50	100	0	0	0	0	15	15	15	75	0	15	0
4	0	0	20	0	35	0	0	50	50	0	50	95	0	0
5	0	0	0	40	35	0	0	0	20	10	50	95	0	35
6	0	0	90	0	35	0	0	0	0	15	0	0	0	0
7	0	20	90	0	0	0	0	0	0	0	20	0	30	0
8	0	0	90	0	15	0	0	0	0	0	0	0	50	0
9	10	90	90	35	50	0	0	0	0	35	0	0	0	25
10	35	80	90	15	25	0	0	0	0	0	0	0	0	35
11	10	80	90	35	0	0	0	50	10	35	0	0	0	0
12	25	80	100	35	0	0	0	0	0	45	15	95	55	0
13	15	40	100	35	0	0	0	50	40	15	20	50	0	0
14	50	80	90	35	0	0	0	0	10	25	35	90	0	35
15	55	90	90	15	50	0	0	0	30	35	0	90	0	35
16	0	80	100	97	0	20	0	45	0	25	35	0	0	0
17	35	50	90	0	0	0	0	0	0	60	35	0	0	0
18	40	0	90	0	0	75	0	0	0	0	60	80	0	30
19	25	15	90	0	0	25	0	0	0	0	0	90	0	0
20	10	15	85	60	55	0	0	0	0	10	15	0	60	35
21	0	65	90	0	70	0	0	0	0	0	0	0	0	0
22	0	50	90	0	100	0	0	0	0	20	50	0	0	25
23	0	15	90	0	0	0	0	0	0	10	0	0	0	0
24	0	85	85	0	0	0	0	0	0	35	0	0	0	0
25	40	90	100	0	0	0	0	0	0	0	0	70	0	0
26	40	100	0	15	0	0	0	0	0	0	0	0	0	0
27	20	85	75	10	0	0	0	0	0	0	0	0	0	0
28	30	90	35	0	10	0	0	0	10	25	15	0	10	0
29	0	0	100	15	30	0	0	50	5	45	5	0	5	55

SUBSTITUTE SHEET

1 The above embodiments and examples are given to
illustrate the scope and spirit of the instant invention.
These embodiments and examples will make apparent, to those
skilled in the art, other embodiments and examples. These
5 other embodiments and examples are within the contemplation
of the present invention. Therefore, this invention should
be limited only by the appended claims.

10

15

20

25

30


35

SUBSTITUTE SHEET

1 WHAT IS CLAIMED IS:

1. A compound having the structural formula

5



10 where R and R¹ are the same or different and are hydrogen or C₁-C₆ alkyl; R² is hydrogen, chlorine or bromine; R³ is hydrogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, trihalomethyl, nitro, cyano, phenyl, phenyl substituted with C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, trihalomethyl, nitro or cyano, phenoxy or phenoxy substituted with C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, trihalomethyl, nitro or cyano; and n is 0 or 1.

20 2. A compound in accordance with Claim 1 wherein R and R¹ are the same or different and are hydrogen or C₁-C₄ alkyl; R² is hydrogen or chlorine; R³ is hydrogen, halogen, phenyl or phenyl substituted with methyl, ethoxy, halogen or trifluoromethyl.

25 3. A compound in accordance with Claim 2 where R and R¹ are the same or different and are hydrogen or C₁-C₃ alkyl; R² is hydrogen; R³ is chlorine, phenyl or phenoxy substituted with halogen or trifluoromethyl.

30 4. A salt comprising the product of the compound of Claim 1 and an organic or inorganic acid.

35 5. A salt in accordance with Claim 4 wherein said acid is hydrochloride acid or methanesulfonic acid.

SUBSTITUTE SHEET

1 6. A composition comprising a fungicidally
effective amount of the compound of Claim 1 and a carrier
therefor.

3 7. A composition comprising a fungicidally
5 effective amount of the salt of Claim 4 and a carrier
therefor.

8. A process for controlling phytopathogenic fungi
10 comprising applying a fungicidally effective amount of the
compound of Claim 1 to the locus under attack by phytopatho-
genic fungi.

15

20

25

30


35

SUBSTITUTE SHEET

INTERNATIONAL SEARCH REPORT

PCT/US 91/07801

International Application No.

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all)		
According to International Patent Classification (IPC) or to both National Classification and IPC Int. Cl. 5 C07D405/12; A01N43/40; //(C07D405/12; 319:00; 213:00) (C07D405/12; 317:00; 213:00)		
II. FIELDS SEARCHED		
Minimum Documentation Searched?		
Classification System	Classification Symbols	
Int. Cl. 5	C07D	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
A	EP,A,0 129 507 (CIBA-GEIGY AG) 27 December 1984 see claims 1,10-14; tables ---	1-8
A	EP,A,0 122 410 (CHUGAI SEIYAKU KABUSHIKI KAISHA) 24 October 1984 see the whole document ---	1-8
A	US,A,4 143 144 (H.K. TOBOL ET AL.) 6 March 1979 cited in the application see claims 1,12; examples ---	1-8
<p>¹⁰ Special categories of cited documents : ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
09 MARCH 1992	17 MAR 1992	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	P. BOSMA 	

Form PCT/ISA/210 (second sheet) (January 1985)

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO. US 9107801
SA 53906**

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 09/03/92

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0129507	27-12-84	AU-B- 571523	21-04-88
		AU-A- 2835984	22-11-84
		CA-A- 1247106	20-12-88
		GB-A, B 2142624	23-01-85
		JP-A- 59222476	14-12-84
		US-A- 4859687	22-08-89
EP-A-0122410	24-10-84	JP-A- 59155360	04-09-84
		JP-A- 59199677	12-11-84
		JP-A- 60136565	20-07-85
		AU-B- 565329	10-09-87
		AU-A- 2448784	30-08-84
		US-A- 4590198	20-05-86
US-A-4143144	06-03-79	AU-B- 519227	19-11-81
		AU-A- 4496479	27-09-79
		CA-A- 1109472	22-09-81
		EP-A, B 0004435	03-10-79

EPO FORM P007

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

